

AD-A162 876 PEAKEDNESS OF WEIGHTED AVERAGES OF JOINTLY DISTRIBUTED 1/1
RANDOM VARIABLES. (U) FLORIDA STATE UNIV TALLAHASSEE
DEPT OF STATISTICS W CHAN ET AL. OCT 85
UNCLASSIFIED FSU-STATISTICS-M712 AFOSR-TR-85-1123 F/G 12/1 NL

PEAKEDNESS OF WEIGHTED AVERAGES OF JOINTLY DISTRIBUTED
RANDOM VARIABLES. (U) FLORIDA STATE UNIV TALLAHASSEE
DEPT OF STATISTICS W CHAN ET AL. OCT 85
FSU-STATISTICS-M712 AFOSR-TR-85-1123 F/G 12/1

141

UNCLASSIFIED

F/G 12/1

NL

END

FILMED

PEAKEDNESS OF WEIGHTED AVERAGES OF
JOINTLY DISTRIBUTED RANDOM VARIABLES

by

Wai Chan¹, Dong Ho Park², and Frank Proschan³,FSU Technical Report No. M712
AFOSR Technical Report No. 85-184

October, 1985

¹Ohio State University
Department of Statistics
Columbus, Ohio 43210

and

²University of Nebraska
Department of Statistics
Lincoln, Nebraska 68588

and

³Florida State University
Department of Statistics
Tallahassee, Florida 32306DTIC
DECEMBER
DEC 31 1985

AD-A162 876

DTIC FILE COPY

Research sponsored by the Air Force Office of Scientific Research under Contract Number ~~AFOSR-85-0007~~. The U.S. Government is authorized to reproduce and distribute reprints for Governmental purposes notwithstanding any copyright notation thereon.

F49620-82-X-0007

Key Words: Peakedness, convex combination, majorization, Schur-concave density, Cauchy distributions.

AMS (1981) subject classification No. 60E15.

85 12 30 015

REPORT DOCUMENTATION PAGE

1. REPORT NUMBER FSU M 712 AFOSA No. 85-185	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and subtitle) Peakedness of Weighted Averages of Jointly Distributed Random Variables		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) Wai Chan, Dong Ho Park, and Frank Proschan		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Florida State University Department of Statistics Tallahassee, Florida 32306 University of Nebraska Ohio State University Department of Statistics Department of Statistics Lincoln, Nebraska Columbus, Ohio 43210 68588		8. CONTRACT OR GRANT NUMBER(s) AFOSR 82-K-0007
		10. PROGRAM ELEMENT, PROJECT, TASK AREA, & WORK UNIT NUMBERS 61102F 2304 A5
		12. REPORT DATE October, 1985
		13. NUMBER OF PAGES 8
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Air Force Air Force Office of Scientific Research Bolling Air Force Base, DC 20332		15. SECURITY CLASS. (of this report) (C)
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
		16. DISTRIBUTION STATEMENT (of this report) Unclassified
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report) Distribution Unlimited		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS Peakedness, Convex combination, majorization, Schur-concave density, Cauchy distributions. Sub A		

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

This note extends the Proschan (1965) result on peakedness comparison for a convex combination of i.i.d. random variables from a PF_2 density. Now the underlying random variables are jointly distributed from a Schur-concave density. The result permits a more refined discription of convergence in the Law of Large Numbers.

Keywords: Cauchy distributions

PEAKEDNESS OF WEIGHTED AVERAGES OF
JOINTLY DISTRIBUTED RANDOM VARIABLES

by

Wai Chan, Dong Ho Park, and Frank Proschan



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability Codes	
Avail and/or	
Dist	Special
A-1	

ABSTRACT

This note extends the Proschan (1965) result on peakedness comparison for a convex combination of i.i.d. random variables from a PF_2 density. Now the underlying random variables are jointly distributed from a Schur-concave density. The result permits a more refined discription of convergence in the Law of Large Numbers.

Copyright

1. Introduction

Proschan (1965) shows that:

1.1 Theorem. Let f be PF_2 , $f(t) = f(-t)$ for all t , X_1, \dots, X_n independently distributed with density f , $p \stackrel{m}{>} p'$, p, p' not identical, $\sum_{i=1}^n p_i = 1 = \sum_{i=1}^n p'_i$. Then $\sum_{i=1}^n p_i X_i$ is strictly more peaked than $\sum_{i=1}^n p'_i X_i$.

(Definitions of majorization ($p \stackrel{m}{>} p'$), PF_2 density, and peakedness are presented in Section 2.) Roughly speaking, Theorem 1.1 states that a weighted average of i.i.d. random variables converges more rapidly in the case in which weights are close together as compared with the case in which the weights are diverse.

In the present note, we extend the basic univariate result to the multivariate situation in which the underlying random variables have a joint Schur-concave density. Theorem 2.3 presents the precise statement of the multivariate extension.

2. Peakedness comparisons

The theory of majorization is exploited in this section to obtain more general versions of the result of Proschan (1965). We begin with some definitions

Definition 2.1. Let $a_1 \geq \dots \geq a_n$ and $b_1 \geq \dots \geq b_n$ be decreasing rearrangements of the components of the vectors a and b . We say that the vector b is majorized by a , written $a \stackrel{m}{\geq} b$ if

$$\sum_{i=1}^n a_i = \sum_{i=1}^n b_i$$

and
$$\sum_{i=1}^k a_i \geq \sum_{i=1}^k b_i \quad \text{for } k=1, \dots, n-1.$$

Definition 2.2. A real valued function f defined on \mathcal{R}^n is said to be a Schur-concave function if $f(\underline{a}) \leq f(\underline{b})$ whenever $\underline{a} \succeq_m \underline{b}$.

A function f defined on \mathcal{R}^n is said to be sign invariant if $f(x_1, \dots, x_n) = f(|x_1|, \dots, |x_n|)$. In the following Lemma, we give a peakedness comparison for random variables with a sign invariant and Schur-concave density.

Theorem 2.3. Suppose the random vector $\underline{X} = (X_1, \dots, X_n)$ has a Schur-concave density f . If f is sign-invariant and satisfies

$$\int_{-\infty}^{\infty} u f(u, u, x_3, \dots, x_n) du < \infty \text{ for all } x_3, \dots, x_n.$$

Then for all $t \geq 0$,

$$\Psi(a_1, \dots, a_n) = P\left(\sum_{i=1}^n a_i X_i \leq t\right)$$

is a Schur-concave function of $\underline{a} = (a_1, \dots, a_n)$, $a_i \geq 0$ for all i . Equivalently,

$\sum b_i X_i$ is more peaked than $\sum a_i X_i$ whenever $\underline{a} \succeq_m \underline{b}$.

Proof.

Without loss of generality, we may assume that $\sum a_i = 1$. We first consider the case $n = 2$.

Let $0 \leq a \leq \frac{1}{2}$ and $\bar{a} = 1 - a$. Let $h(a) = P(\bar{a}X_1 + aX_2 \leq t) = \int_{-\infty}^{\infty} G_{X_2|X_1=u}\left(\frac{t-au}{\bar{a}}\right) g_1(u) du$ where g_1 is the marginal density of X_1 and $G_{X_2|X_1=u}$ is the conditional distribution function of X_2 given that $X_1 = u$.

Differentiation under the integral sign is permissible here, so that

$$\begin{aligned} \bar{a}^2 h'(a) &= \int_{-\infty}^{\infty} g_{X_2|X_1=u}(\frac{t-au}{\bar{a}}) g_1(u) (t-u) du \\ &= \int_{-\infty}^{\infty} f(u, \frac{t-au}{\bar{a}}) (t-u) du. \\ &= \int_{-\infty}^t f(u, \frac{t-au}{\bar{a}}) (t-u) du \\ &\quad + \int_t^{\infty} f(u, \frac{t-au}{\bar{a}}) (t-u) du. \end{aligned}$$

Now let $v = t - u$ in the first integral and $v = u - t$ in the second integral. We obtain

$$\begin{aligned} \bar{a}^2 h'(a) &= \int_0^{\infty} v [f(t-v, t+\frac{a}{\bar{a}}v) - f(t+v, t-\frac{a}{\bar{a}}v)] dv \\ &= \int_0^{\infty} v [f(v-t, \frac{a}{\bar{a}}v+t) - f(v+t, \frac{a}{\bar{a}}v-t)] dv, \end{aligned}$$

since f is sign invariant. But this is nonpositive because

$$(v+t, \frac{a}{\bar{a}}v-t) \stackrel{m}{\geq} (v-t, \frac{a}{\bar{a}}v+t)$$

and f is Schur-concave. Thus $h(a)$ is increasing in a , $0 \leq a \leq \frac{1}{2}$.

The result for $n \geq 3$ now follows since

$$\begin{aligned} P(\sum a_i X_i \leq t) \\ = E [P(a_1 X_1 + a_2 X_2 \leq t - \sum_{i=3}^n a_i X_i | X_3, \dots, X_n)] \end{aligned}$$

and the conditional density $f(x_1, x_2 | x_3, \dots, x_n)$ is also Schur-concave and sign invariant. \square

Remark 2.4. To justify differentiation under the integral sign, we note that

$$\begin{aligned} &\int_{-\infty}^{\infty} |f(u, \frac{t-au}{\bar{a}}) (t-u)| du \\ &\leq \int_{-\infty}^{\infty} |t-u| f(\frac{|u-t|}{\bar{a}}, \frac{|u-t|}{\bar{a}}) du < \infty, \end{aligned}$$

which follows from (2.1).

This condition is clearly not a necessary condition, but it can be easily verified for most Schur-concave multivariate distributions. For example, the multivariate Cauchy density:

$$f(x_1, \dots, x_n) = \pi^{-(n+1)/2} \Gamma((n+1)/2) \left(1 + \sum_{i=1}^n x_i^2\right)^{-(n+1)/2}$$

has this property.

The following result is an immediate application of Theorem 2.3.

Corollary 2.5. Let X_1, \dots, X_n be random variables with joint Schur-concave density f . Let f be sign invariant and satisfy

$$\int_{-\infty}^{\infty} u f(u, u, x_2, \dots, x_n) du < \infty \text{ for all } x_2, \dots, x_n.$$

Then $\frac{1}{k} \sum_{i=1}^k X_i$ is increasing in peakedness as k increases from 1 to n .

Proof.

Let $\underline{a}_1 = (1, 0, \dots, 0)$, $\underline{a}_2 = (\frac{1}{2}, \frac{1}{2}, 0, \dots, 0), \dots$ and $\underline{a}_n = (\frac{1}{n}, \dots, \frac{1}{n})$ where each vector contains n components. Then $\underline{a}_1 \geq \underline{a}_2 \geq \dots \geq \underline{a}_n$. The result follows from Theorem 2.3. \square

Suppose $\underline{X} = (X_1, \dots, X_n)$ and $\underline{Y} = (Y_1, \dots, Y_n)$ are independently distributed with respective densities f and g where both f and g are Schur-concave and sign invariant, Theorem 2.3 implies that $\sum b_i(X_i + Y_i)$ is more peaked than $\sum a_i(X_i + Y_i)$ whenever $\underline{a} \geq \underline{b}$. This is true because the convolution of Schur-concave functions is Schur-concave. However, if Y_1, \dots, Y_n are i.i.d. Cauchy, then the joint density given by

$$g(X_1, \dots, X_n) = \left(\frac{a}{\pi}\right)^n \prod_{i=1}^n (1 + a^2 X_i^2)^{-1}, \quad a > 0,$$

is not Schur-concave. Theorem 2.7 below, we give conditions on f for which (2.2) holds. First we prove the following Lemma.

Lemma 2.6. Let $\underline{X} = (X_1, \dots, X_n)$ and $\underline{Y} = (Y_1, \dots, Y_n)$ be independently distributed with respective densities f_1 and f_2 . Suppose $f_i(t_1, \dots, t_n)$ is symmetric with respect to zero and nonincreasing in each argument for $t_k > 0$, $k = 1, \dots, n$, and for $i = 1, 2$.

Let $\sum_{i=1}^n b_i X_i$ be more peaked than $\sum_{i=1}^n a_i X_i$ and $\sum_{i=1}^n b_i Y_i$ be more peaked than $\sum_{i=1}^n a_i Y_i$ where $a_i \geq 0$ and $b_i \geq 0$ for

$i = 1, \dots, n$. Then

$$\sum_{i=1}^n b_i (X_i + Y_i) \text{ is more peaked than } \sum_{i=1}^n a_i (X_i + Y_i).$$

Proof.

This result follows immediately from the Lemma of Birnbaum (1948) by noting that the random variables $\sum_{i=1}^n a_i X_i$, $\sum_{i=1}^n a_i Y_i$, $\sum_{i=1}^n b_i X_i$, and $\sum_{i=1}^n b_i Y_i$ have symmetric and unimodal densities. \square

The following theorem identifies a different class of densities for which the conclusion of Theorem 2.3 holds.

Theorem 2.7. Suppose that the random vector $\underline{X} = (X_1, \dots, X_n)$ has a Schur-concave sign-invariant density f . Let f be nonincreasing in each argument over the positive values and satisfy (2.1). Let Y_1, \dots, Y_n be i.i.d. Cauchy with joint density g as given in (2.3). Let \underline{X} and $\underline{Y} = (Y_1, \dots, Y_n)$ be independent, and $\underline{a} \succeq \underline{b}$ where

$a_i \geq 0$, $b_i \geq 0$ for all i and $1 = \sum_{i=1}^n a_i = \sum_{i=1}^n b_i$. Then

$$\sum_{i=1}^n b_i (X_i + Y_i) \text{ is more peaked than } \sum_{i=1}^n a_i (X_i + Y_i).$$

Proof.

We use the fact that $\sum_{i=1}^n a_i Y_i$, $\sum_{i=1}^n b_i Y_i$ have the same distribution as does Y_1 .

From Theorem 2.3, $\sum_{i=1}^n b_i X_i$ is more peaked than is $\sum_{i=1}^n a_i X_i$. The result now follows from Lemma 2.6. \square

REFERENCES

- [1] Birnbaum, Z. W. (1948). On random variables with comparable peakedness.
Ann. Math. Statist. 19. 76-81.
- [2] Proschan, F. (1965). Peakedness of distributions of convex combinations.
Ann. Math. Statist. 36. 1703-1706.

END

FILMED

2-86

DTIC